

General Procedure Instructions

Read and follow these instructions closely. Failure to follow them will result in your procedure being **cancelled or rescheduled** to a different day. These instructions are for your safety. Failure to follow them could cause severe complications, resulting in **HOSPITALIZATION or even DEATH**.

****DO NOT DRINK ANYTHING FOR FOUR (4) HOURS BEFORE YOUR SCHEDULED PROCEDURE START TIME. THIS EVEN INCLUDES CHEWING GUM, MINTS, OR SMALL AMOUNTS OF WATER AND MEDICATIONS.****

****NO TOBACCO PRODUCTS SIX (6) HOURS BEFORE YOUR SCHEDULED PROCEDURE START TIME.****

****No Illicit Drugs (Meth, Cocaine, Heroin, etc.) 7 days prior to your procedure****

****Avoid CBD Oil and Marijuana use 24 hours prior to your procedure.****

****Please notify us** if you are taking any **blood thinning** medications, (ex: Coumadin, Plavix, Aggrenox, Xarelto, Eliquis, Pletal, Pradaxa, etc.) or Diabetic Medication.

Do NOT take any of the following medications for **SEVEN (7)** days prior to your procedure:

1. Iron medications or supplements (e.g. Fermalox, Niferex, Ferrous Sulfate), or vitamins containing iron
2. Any fiber supplements such as Metamucil, Citrucel, Konsyl

DO:

1. Call the facility where you will have your procedure with any questions regarding your medications or preparation instructions.
2. **Have an adult (18+) accompany you to the procedure. He/she MUST stay at the center the entire time you are there to be available for any emergencies and because some of the medication you will receive will impair your ability to drive.**
3. If you are having a colonoscopy, follow the prep instructions carefully and completely.
4. If you are having an upper endoscopy (EGD), begin a clear liquid diet at 11:00 PM the night before your procedure.
5. Stay hydrated by drinking plenty of clear liquids up to your four hour cutoff point.
6. Brush your teeth, but do not swallow water or toothpaste.
7. Check with the physician managing your medications regarding options for temporarily stopping any blood thinning medications or diabetic medications.
8. Take your normal dose of blood pressure, heart, or seizure medications. If, you should take medication the morning of your procedure, take it **at least four (4) hours prior to your procedure**.
9. Arrive for your procedure as directed. Your arrival time will be approximately one hour before your procedure begins. However, if your procedure is scheduled for 7:00am or 7:30am, please be aware that our office does not open until 6:30am. You do not need to arrive before 6:30am.
10. Because each case is different, many factors determine what time your procedure will actually begin. Please remain patient. We will do our best to accommodate you in a timely manner.



Digestive Care Center

Rectal Endoscopic Ultrasound Prep

*Because of the medications you will receive, ***you must have someone accompany you that can stay in the Endoscopy Center waiting room while you have your procedure and drive you home after the procedure.***

Please bring with you any x-ray films that may have been taken relating to this appointment.

Preparing for your Endoscopic Ultrasound:

Please notify us if you are taking any **blood thinning** medications, (ex: Coumadin, Xarelto, Eliquis, Pradaxa, etc). **Refer to appointment information for specific stop times regarding medication you take regularly.**

ALWAYS check with the physician managing your medication before stopping any blood thinning or diabetic medications.

No solid foods after midnight. You may drink clear liquids up to **4 hours** before your procedure, (examples – Fat free broth or bouillon, plain gelatin desserts, gingerale, ice tea, coffee, kool-aid, clear fruit juices, Hi-C, carbonated beverages, (i.e. Coke, Sprite, Pepsi). Please avoid liquids containing red, purple or orange dye. **No** milk, orange or tomato juice.

You must take any heart, blood pressure or seizure medication that you normally take with a **small sip** of water the morning of your procedure.

If you are scheduled for a rectal ultrasound, you will need to give yourself 2 enemas one hour before leaving home – insert one and hold then give the other and release.

*****Please note: Some insurances do require outpatient precertification. It is your responsibility to notify us if precert is required for your procedure.***

IF YOU HAVE ANY QUESTIONS, PLEASE CALL 812-477-6103

Thank You